

Live vaccine for
the first shot
in puppies
at the age
six weeks



Biocan DHPPi

lyophilisate for preparation of solution for injection

Live vaccine against canine distemper (CDV), inf. laryngotracheitis (CAV-2), inf. hepatitis (CAV-1), parvovirus (CPV-2), parainfluenza (CPIV-2) in dogs.

COMPOSITION

Composition – 1 ml:

Freeze-dried component:

Virus febris contagiosae canis

min. $10^{3.0}$ TCID₅₀, max. $10^{4.5}$ TCID₅₀

Virus laryngotracheitidis

contagiosae canis min. $10^{3.5}$
TCID₅₀, max. $10^{4.5}$ TCID₅₀

Parvovirus enteritidis canis

min. $10^{4.5}$ TCID₅₀, max. $10^{5.5}$ TCID₅₀

Virus parainfluenzae canis

min. $10^{3.0}$ TCID₅₀, max. $10^{4.2}$ TCID₅₀

TARGET SPECIES

Dogs.

INDICATION

For the active immunization of dogs against distemper, infectious hepatitis, infectious laryngotracheitis, parvovirus and parainfluenza.

INTERACTION

The vaccine Biocan DHPPi can be used separately or simultaneously with other vaccines Biocan according to recommended vaccination scheme or in group with fluid vaccines Biocan (LR, L, C, R).

DOSAGE

The dose is 1 ml regardless of age, weight and breed of the individual, vaccination may be first performed during the sixth week of age.

Method of administration – subcutaneously, preferably in zone behind the shoulder-blade.

SHELF LIFE

24 months. When diluted, the vaccine shall be used immediately.

STORAGE

Store in a dry and dark place at the temperature of 2 °C – 8 °C.

PACKAGE

5 × 1 ml of vaccine Biocan DHPPi

+ 5 × 1 ml of diluent

10 × 1 ml of vaccine Biocan DHPPi + 10 × 1 ml of diluent

50 × 1 ml of vaccine Biocan DHPPi + 50 × 1 ml of diluent