

Product for  
treatment acute  
and chronic  
mastitis during  
lactation with the  
unique antibiotic  
Novobiocinum



# GAMARET

## intramammary suspension

### COMPOSITION

1 applicator 10 ml contains:  
*Procaini benzylpenicillinum*  
*monohydricum* 100 mg  
*Neomycini sulfas* 102 000 IU  
*Dihydrostreptomycini sulfas* 91 250 IU  
*Novobiocinum natriicum* 100 mg  
*Prednisolonum* 10 mg

### TARGET SPECIES

Cattle – milking cows in lactation period.

### INDICATIONS

Treatment of acute and chronic mastitis in cows in the period of lactation caused by microorganisms susceptible to novobiocin, penicillin, dihydrostreptomycin and neomycin.

### DOSAGE

The content of one applicator (10 ml of the preparation) is administered to the affected quadrant. Before the preparation is administered the lacteal glands are milked and thoroughly cleaned and disinfected using enclosed cleaning wipes.

#### Administration method:

The preparation is intended only for intramammary use. Shake well before use. Udder and teats are washed as required with warm water and thoroughly dried. The ends of teats are disinfected with an appropriate agent and after removing the plastic cover of the applicator's tip is the applicator inserted to the teat duct. The content is administered to the udder by pressing the piston. After the administration is the particular quadrant massaged in order to ensure a better distribution of the preparation into milk cisterns. If required by the situation, it is possible to repeat therapy after 24–48 hours.

### WITHDRAWAL PERIODS

Meat 7 days, milk – 72 hours (6 milking).

### SHELF LIFE

Shelf-life in an intact package: 18 months

### STORAGE

Store below 25 °C and keep in dry place and protect for light. Do not freeze.

### PACKAGE

10 ml plastic applicators (packaging 20 applicators in paper box).